



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

1. *An Anatomical Account of some Remarkable things, found on the Dissection of a Woman, who dyed of the Dropsie, after the Paracentesis was performed, with a small Reflection on the Causes of the Dropsie. By Dr. Charles Preston.*

Being called to the Dissection of Madam *Vaillant*, the Chirurgion one *Du Linier*, Operator to Monsieur *Du Vergy*; what we found remarkable is as follows :

In opening the Lower Belly, the Liver was found very White, the Epiploon extreamly dried, the Stomach much bigger than ordinary, the winding of the Colon, which passes under the Stomach, extreamly drawn together by three Threds. In the Umbilical Region, the Intestines Jejunum and Ilium, much inflamed, and their Tunicks much more thick than ordinary. In the Hypogastrick Region, all the inferiour part of the Intestine Ileum, on that side near the Bladder, and all the bottom of the Matrix; as also the inferiour part of the Rectum much Inflamed and Ulcerated. In the bottom of the Matrix there was an Abscess, and the internal Orifice extreamly Dilated, about the largeness of a Crown; the extremity of the inferiour part of the Ureter Cartilaginous; the Extremity of the *Tubæ Fallopianæ* went so high, as the second Vertebre of the Lumbar Region; in the interior part it was Dilated six Lines, and near the bottom of the Matrix, about two Inches, and was tyed to all the inferior part of the Kidney; that of the left side was Dilated about four Lines in the upper part, and six in the inferior.

The right Testicle, or Ovarium, which ordinarily does not exceed the bigness of a Pidgeons Egg, was here of a considerable bigness, three Inches long, and two of breadth; and in the inferior part, there was found an Egg, hanging by its Ligament, out of the *Tuba Fallopiana*, about the bigness of the Yolk of a common Hens Egg;

Egg; which, for Experiment, I caused to be Boiled, and it hardned as an ordinary Egg, but did not pursue the Tryal any farther.

The right Kidney was of a considerable bigness, went up as far as the last of the true Ribs, and descended below the Umbilical Region; the Pelvis was Dilated about three Inches in breadth, and seven in length.

In the second Cavity, the Lungs were of a Livid Colour, as in all Chronical Diseases; and on the right side were adherent to the Membrane Pleura; and on the left side was an adherence of the Inferior Lobe to the Diaphragm: In the Pericard was little or no Serum; and what we found of a bloody Colour in dissecting the Heart, we found a great Polypus in the right Ventricle, taking up almost all the Cavity, about five or six Lines in thickness, and half a Foot in length.

Lastly, It is to be Noted, that there was not much Water, there having a great deal run out in the Operation.

Having given a Relation of what was found most Remarkable, on the Dissection of this Hydropick Woman, I shall only make a short Reflection, as to the Causes of the Dropsie; for there is here such a complex of Diseases, as would furnish Matter for a Volume, but I reckon most of them effects of the Dropsie.

We find some Authors blame the Liver, because it is found of a whitish Colour; others the Spleen, because sometimes Schirrous; but these are only the effects of the Disease, and the present Subject furnishes me with Arguments; for altho' the Liver was of a white Colour without, yet its interior was Red. *Sylvius* has observed a Rupture of the Lacteal Veins, the *Receptaculum Chyli*, or *Ductus Thoracicus*; but because that is reckoned incurable I shall pass to others. Since the Discovery of the lymphatick Vessels, they have not stood to say, that a Rupture of them is the cause; but Anatomy tells us they are not so easily broken; and upon the dissection of hydropick Bodies,

you

you will never find any of the lymphatick Vessels broken.

We shall then take the true Cause from the Mechanick Structure of the parts, and the disposition of the Blood, which are first the Relaxation of the Fibres and Pores of the Vessels, or the *Vesiculæ*, which are between the Arteries and the Veins, or secondly, a compression of the Vessels; for the Lymphaticks take their Origine from the Membranes, which covers the Muscles *Viscera* and *Glands*; therefore, when the *Vesiculæ* are too much straitned with Serosity, their Fibres loose their natural force, and become incapable to expel the too great quantity of Water; but the *Vesiculæ* are enlarged from day to day, until their Fibres suffer so great an Extension even as to break, from hence is the source of those Waters; it happens also sometimes, that the pores of the said *Vesiculæ* are so widened, that the Lymph runs into the Cavity of the Belly, or the Interstices of the Muscles.

The Cause from the Disposition of the Blood, is either when it is too thin or too viscid; too thin, that it passes easily through the Pores of the Vesicles; too viscid, that it cannot pass through the Capillary Vessels, and by consequence Compresses the adjacent parts, so causes Obstructions. I shall insist no further as to the Cause, only take Notice, whether in reason this Woman could have lived any time longer, considering all the Complex of Diseases; or if the Operation of the Paracentesis was the Cause of her being sooner carried off: From what is said above, it is impossible she could Recover; only this is to be remarked, that where the Dropsie is of a long continuance, and the Persons much Debilitated and of Age, in that Case the Operation is not to be performed; for generally the *Viscera* are Corrupted.

When ye perform the Operation, Extract the Water by degrees, and not all at once, else you endanger the Person, for scarce One escapes of a Hundred that is done otherwise.

II. *An*